


9-1-2016

Rapid Museum

Gary Barwin

Niagara Regional Rhyme Gland Laboratory

Follow this and additional works at: <http://scholars.wlu.ca/thegoose>

 Part of the [Biology Commons](#), [Critical and Cultural Studies Commons](#), [Entomology Commons](#), [Literature in English, North America Commons](#), [Nature and Society Relations Commons](#), [Place and Environment Commons](#), [Poetry Commons](#), and the [Population Biology Commons](#)

Recommended Citation

Barwin, Gary (2016) "Rapid Museum," *The Goose*: Vol. 15: Iss. 1, Article 41.

Available at: <http://scholars.wlu.ca/thegoose/vol15/iss1/41>

This Poetry is brought to you for free and open access by Scholars Commons @ Laurier. It has been accepted for inclusion in The Goose by an authorized administrator of Scholars Commons @ Laurier. For more information, please contact scholarscommons@wlu.ca.

Rapid Museum

hasings model G regressed on V as performed using G. srbeser of bee-
 ches, collected in the same decade at a more or less fixed size
 2W. Joid the veral change in shape, if, ob being affected by time
 are external factors! Analysis were performed on: Statistica 6.1,
 tory, t-test, and ANOVA with Excel, 2007. Furthermore, mid-
 latitude land precipitation shows a 0.5–1% increase per decade
 while sub-tropical latitudes show a 0.3% decrease per decade
 (Figs 1, 2001). More research is required at the regional level to
 reflect this heterogeneity, and we know of no studies that have
 examined changes in G. minima from 1980 to 1989, increased by
 0.8% per decade observed linear regression $\ln(\text{mean } \mu) =$
 $\ln(0.14) + 0.008 \times \text{Decade}$, Fig. 2A). The fineness ratio (length/width)
 increased by 3% based on a linear regression $\ln(\text{width}) =$
 $\ln(0.09) + 0.003 \times \text{Decade}$ (Fig. 2B). The biometric and geographic
 variables examined also changed over time: Mean annual tem-
 peratures are from southern U.S.A., an area which is known

hasings model G regressed on V as performed using G. srbeser of bee-
 ches, collected in the same decade at a more or less fixed size
 2W. Joid the veral change in shape, if, ob being affected by time
 are external factors! Analysis were performed on: Statistica 6.1,
 tory, t-test, and ANOVA with Excel, 2007. Furthermore, mid-
 latitude land precipitation shows a 0.5–1% increase per decade
 while sub-tropical latitudes show a 0.3% decrease per decade
 (Figs 1, 2001). More research is required at the regional level to
 reflect this heterogeneity, and we know of no studies that have
 examined changes in G. minima from 1980 to 1989, increased by
 0.8% per decade observed linear regression $\ln(\text{mean } \mu) =$
 $\ln(0.14) + 0.008 \times \text{Decade}$, Fig. 2A). The fineness ratio (length/width)
 increased by 3% based on a linear regression $\ln(\text{width}) =$
 $\ln(0.09) + 0.003 \times \text{Decade}$ (Fig. 2B). The biometric and geographic
 variables examined also changed over time: Mean annual tem-
 peratures are from southern U.S.A., an area which is known

hasings model G regressed on V as performed using G. srbeser of bee-
 ches, collected in the same decade at a more or less fixed size
 2W. Joid the veral change in shape, if, ob being affected by time
 are external factors! Analysis were performed on: Statistica 6.1,
 tory, t-test, and ANOVA with Excel, 2007. Furthermore, mid-
 latitude land precipitation shows a 0.5–1% increase per decade
 while sub-tropical latitudes show a 0.3% decrease per decade
 (Figs 1, 2001). More research is required at the regional level to
 reflect this heterogeneity, and we know of no studies that have
 examined changes in G. minima from 1980 to 1989, increased by
 0.8% per decade observed linear regression $\ln(\text{mean } \mu) =$
 $\ln(0.14) + 0.008 \times \text{Decade}$, Fig. 2A). The fineness ratio (length/width)
 increased by 3% based on a linear regression $\ln(\text{width}) =$
 $\ln(0.09) + 0.003 \times \text{Decade}$ (Fig. 2B). The biometric and geographic
 variables examined also changed over time: Mean annual tem-
 peratures are from southern U.S.A., an area which is known

Note on the Text

How has human communication with the earth (in the form of climate change) affected its language (the shape of stream beetles) and specifically its lexicon (in this case, samples of stream beetles found in museums)? How quickly is this language changing? How urgently is it sending communications back to us?

Rapid Museum is a visual poem based on a scientific (ecology/biology) paper by Jennifer Babin-Fenske, Madhur Anand and Yves Alarie: "Rapid morphological change in stream beetle museum specimens correlates with climate change." Madhur Anand's poetry collection, *A New Index for Predicting Catastrophes* (McClelland and Stewart, 2015), uses some textual material directly taken from her own scientific articles, and she invited me to similarly take text from one of her articles and explore how it might be used in a poem. I created a visual poem but also a video poem (<http://bit.ly/2adz9d5>) incorporating the visual poem and audio derived from the text of the article.

Thanks to Madhur Anand for inviting me to explore the artistic possibilities of her scientific texts.

Works Cited

Babin-Fenske, J., Anand, M. and Alarie, Y. (2008), Rapid morphological change in stream beetle museum specimens correlates with climate change. *Ecological Entomology*, 33: 646–651. doi: 10.1111/j.1365-2311.2008.01018.x

GARY BARWIN is a writer, composer, and multimedia artist and the author of 20 books of poetry and fiction, including the recent national bestselling novel, *Yiddish for Pirates. No TV for Woodpeckers*, a new poetry collection, will appear from Wolsak & Wynn in 2017. Barwin was writer-in-residence at Western University last year and currently teaches at Mohawk College and is writer-in-residence in several youth shelters and custody facilities in Hamilton, Ontario.