

Wilfrid Laurier University

## Scholars Commons @ Laurier

---

Physics and Computer Science Faculty  
Publications

Physics and Computer Science

---

1974

### A Note on Chambers' Method

James A. Blackburn

Wilfrid Laurier University, jabjabjab@cogeco.ca

Y. Beaudoin

University of Western Ontario

Follow this and additional works at: [https://scholars.wlu.ca/phys\\_faculty](https://scholars.wlu.ca/phys_faculty)

---

#### Recommended Citation

Blackburn, James A. and Beaudoin, Y., "A Note on Chambers' Method" (1974). *Physics and Computer Science Faculty Publications*. 32.

[https://scholars.wlu.ca/phys\\_faculty/32](https://scholars.wlu.ca/phys_faculty/32)

This Article is brought to you for free and open access by the Physics and Computer Science at Scholars Commons @ Laurier. It has been accepted for inclusion in Physics and Computer Science Faculty Publications by an authorized administrator of Scholars Commons @ Laurier. For more information, please contact [scholarscommons@wlu.ca](mailto:scholarscommons@wlu.ca).

## A Note on Chambers' Method

By J. A. Blackburn\* and Y. Beaudoin

**Abstract.** A correction is given for one of Chambers' second-order iteration formulae. It is shown that composition of the secant method with itself exhibits a convergence exponent of 2.414, whereas composition of the iteration function with itself yields an exponent of 2.831.

The improved second-order iteration schemes presented recently by L. G. Chambers [1] yield estimates of isolated zeros of  $f(x)$  which exhibit superquadratic convergence exponents. We wish to point out an error in that paper: namely that Eq. (2.10) which was given as

$$X_{n+1} = \frac{X_{n-1} Y_n^2}{(Y_{n-1} - Y_n)^2} + \left( \frac{Y_n}{Y_n - Y_{n-1}} \right) \cdot \left( \frac{Y_n - X_n Y_n'}{Y_n'} \right)$$

should have read

$$(1) \quad X_{n+1} = \frac{Y_n(X_{n-1} Y_n - X_n Y_{n-1})}{(Y_n - Y_{n-1})^2} + \left( \frac{Y_{n-1}}{Y_n - Y_{n-1}} \right) \cdot \left( \frac{Y_n - X_n Y_n'}{Y_n'} \right).$$

The convergence exponent of 2.414 remains valid however.

In a follow-up note on Chambers' method, M. G. Cox [2] pointed out that "an iteration function constructed by composition of the secant method with itself also requires just two function evaluations per iteration and has a convergence exponent of  $(1.618)^2 = 2.618$ ". We believe this remark to be in error for the reasons given below.

Composition of the secant method with itself would generate the two-stage iteration:  $\phi(X_{n-1}, X_n) = X_n^*$ ;  $\phi(X_n, X_n^*) = X_{n+1}$ . The first stage has an associated error [3] of  $\epsilon_n^* = K_0 \epsilon_{n-1} \epsilon_n$  while the second stage is governed by  $\epsilon_{n+1} = K_1 \epsilon_n \epsilon_n^*$ . Therefore

$$(2) \quad \epsilon_{n+1} = K \epsilon_{n-1} \epsilon_n^2.$$

Note that this recurrence relation is of the very same form as that given by Chambers in connection with iteration formulae of the Wegstein type when  $g = 1$ ; the convergence exponent in such a case was 2.414. It is incorrect to merely square the convergence exponent of a single secant method step since this would ignore the role of  $X_n^*$ . Cox's figure of 2.618 is thus overly optimistic although we concede that the practical significance of such a small difference would probably be nil.

Cox also states that composition of the iteration function with itself yields "a convergence exponent of  $(1.839)^2 = 3.382$ , which is superior to all those derived

---

Received May 15, 1973.

AMS (MOS) subject classifications (1970). Primary 65H05; Secondary 65B99.

Key words and phrases. Numerical analysis, approximation to the root of an equation.

\* Research supported by the National Research Council of Canada.

Copyright © 1974, American Mathematical Society

by Chambers". This method would involve the two-stage procedure:  $\phi(X_{n-2}, X_{n-1}, X_n) = X_n^*$ ;  $\phi(X_{n-1}, X_n, X_n^*) = X_{n+1}$ . Two new function evaluations per iteration are required—that is  $f(X_n)$  and  $f(X_n^*)$ . The first and second stage errors are then  $\epsilon_n^* = K_0 \epsilon_{n-2} \epsilon_{n-1} \epsilon_n$  and  $\epsilon_{n+1} = K_1 \epsilon_{n-1} \epsilon_n \epsilon_n^*$ . Therefore,

$$(3) \quad \epsilon_{n+1} = K \epsilon_{n-2}^2 \epsilon_{n-1}^2 \epsilon_n^2.$$

Following Chambers, we assume  $\epsilon_{n+1} = M \epsilon_n^\mu$  and so  $\mu = 1/\mu^2 + 2/\mu + 2$ , the appropriate root of which is 2.831. We conclude therefore that the convergence exponent arising, when the iteration function is cycled on itself, is 2.831 and not 3.382 as given by Cox. Hence, supercubic performance does not occur.

The relative merits of the various methods are clearly revealed by the convergence rates *per function evaluation* [4]: secant  $[1.618]$ ; selfcomposed secant  $[(2.414)^{1/2} = 1.554]$ ; quadratic inverse interpolation  $[1.839]$ ; selfcomposed quadratic inverse interpolation  $[(2.831)^{1/2} = 1.683]$ ; Chambers' first method  $[2^{1/2} = 1.414]$ ; Chambers' second method  $[(2.414)^{1/2} = 1.554]$ ; Chambers' third method  $[(2.732)^{1/2} = 1.653]$ .

Physics Department  
Wilfrid Laurier University  
Waterloo, Ontario, Canada

Department of Applied Mathematics  
University of Western Ontario  
London, Ontario, Canada

1. LL. G. CHAMBERS, "A quadratic formula for finding the root of an equation," *Math. Comp.*, v. 25, 1971, pp. 305–307. MR 45 #4625.

2. M. G. COX, "A note on Chambers' method for finding a zero of a function," *Math. Comp.*, v. 26, 1972, p. 749.

3. D. M. YOUNG & R. T. GREGORY, *A Survey of Numerical Mathematics*, Vol. I, Addison-Wesley, Reading, Mass., 1972, pp. 150–153.

4. We are indebted to the referee for suggesting this final comparison.